

Table S1. Predictors of non-communicable disease behavioral risk factors among adolescents: unadjusted ORs (additional information for Table 4)

Variable	Inadequate FV intake		Unhealthy diet (junk)		SSB		Physical inactivity		Tobacco		Alcohol	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Gender												
Boy	0.89 (0.65–1.22)	0.48	1.01 (0.74–1.39)	0.939	0.95 (0.74–1.23)	0.712	1.09 (0.79–1.49)	0.609	1.17 (0.77–1.79)	0.467	1.06 (0.69–1.63)	0.796
Girl (ref.)	1		1		1		1		1		1	
School location												
Urban area	0.07 (0.04–0.12)	0.001	4.56 (3.14–6.63)	0.001	2.37 (1.83–3.06)	0.001	13.60 (8.21–22.5)	0.001	0.46 (0.29–0.72)	0.001	0.49 (0.31–0.78)	0.002
Rural area (ref.)	1		1		1		1		1		1	
Age (y)												
≤14	1.75 (1.28–2.39)	0.001	0.97 (0.70–1.33)	0.826	0.96 (0.75–1.23)	0.75	0.87 (0.63–1.19)	0.37	1.44 (0.94–2.19)	0.094	0.62 (0.48–0.98)	0.039
>14 (ref.)	1		1		1		1		1		1	
Father's education												
Primary	0.60 (0.32–1.12)	0.106	2.07 (1.19–3.60)	0.01	1.93 (1.23–3.03)	0.004	1.45 (0.78–2.75)	0.241	0.66 (0.34–1.26)	0.209	0.63 (0.28–1.38)	0.245
Secondary	1.24 (0.75–2.05)	0.41	1.73 (1.15–2.75)	0.019	1.63 (1.10–2.47)	0.014	0.85 (0.51–1.43)	0.543	0.39 (0.22–0.71)	0.002	0.89 (0.46–1.71)	0.73
Diploma/degree	0.90 (0.35–2.27)	0.815	0.79 (0.37–1.71)	0.555	1.11 (0.56–2.21)	0.773	0.48 (0.21–1.07)	0.072	1.26 (0.51–3.14)	0.618	0.43 (0.09–2.00)	0.282
No formal education (ref.)	1		1		1		1		1		1	
Mother's education												
Primary	0.70 (0.43–1.14)	0.148	1.13 (0.75–1.71)	0.549	1.82 (1.29–2.58)	0.001	1.20 (0.77–1.89)	0.427	0.76 (0.43–1.34)	0.339	0.99 (0.56–1.76)	0.975
Secondary	2.10 (1.49–2.97)	0.001	1.37 (0.95–1.99)	0.094	1.14 (0.86–1.51)	0.358	0.64 (0.45–0.90)	0.011	0.70 (0.42–1.15)	0.16	1.08 (0.66–1.76)	0.763
Diploma/degree	6.00 (1.69–21.28)	0.006	0.53 (0.13–2.09)	0.363	0.61 (0.17–2.13)	0.437	0.18 (0.05–0.63)	0.008	6.23 (1.76–22.90)	0.006	1.28 (0.16–10.40)	0.817
No formal education (ref.)	1		1		1		1		1		1	
Father's occupation												
Skilled	0.59 (0.18–3.53)	0.562	1.86 (0.31–11.20)	0.5	1.24 (0.21–7.33)	0.815	3.71 (0.68–20.50)	0.132	0.24 (0.04–1.54)	0.132	0	
Semiskilled	0.69 (0.13–3.83)	0.673	2.02 (0.36–11.20)	0.422	0.79 (0.14–4.36)	0.786	3.55 (0.71–17.90)	0.124	0.21 (0.04–1.16)	0.073	0	
Unskilled	0.26 (0.05–1.42)	0.119	3.11 (0.56–17.30)	0.194	1.10 (0.29–6.04)	0.916	6.60 (1.31–33.30)	0.022	0.16 (0.03–0.91)	0.039	0	
No employment (ref.)	1		1		1		1		1		1	
Mother's occupation												
Non-working	0.99 (0.69–1.44)	0.968	1.03 (0.70–1.57)	0.895	0.77 (0.56–1.04)	0.088	1.21 (0.84–1.74)	0.308	0.94 (0.57–1.55)	0.818	1.02 (0.60–1.70)	0.956
Working (ref.)	1		1		1		1		1		1	

The table presents different behavioral patterns among school adolescents. It includes OR, 95% CI, and p-values across various demographic categories such as gender, school location, age, fathers' and mothers' education, and parental occupation. A p-value less than 0.05 is typically considered statistically significant.

OR, odds ratio; FV, fruit and vegetable; CI, confidence interval; SSB, sugar-sweetened beverage; ref., reference category.